25

30

1. A pharmaceutical composition having therapeutic synergy comprising:

What is claimed is:

20(S)-camptothecin, an analog of 20(S)-camptothecin, or a derivative of 20(S)-camptothecin; and

at least one of an alkylating agent, epidophyllotoxin, antimetabolite, antibiotic or vinca alkaloid.

- 2. A pharmaceutical composition according to claim 1, wherein the alkylating agent is cyclophosphamide, ifosfamide, melphalan, hexamethylmelamine, thiotepa or dacarbazine.
- 3. A pharmaceutical composition according to claim 1, wherein the antimetabolite is 5-fluorouracil, cytarabine or a folic acid analog.
- 4. A pharmaceutical composition according to claim 1, wherein the antibiotic is daunorubicine, doxorubicin, bleomycin or mitomycin.
- 5. A pharmaceutical composition according to claim 1, wherein the vinca alkaloid is vinblastine, vincristine, or their synthetic analogues.
- 6. A pharmaceutical composition according to claim 1, wherein the epidophyllotoxin is etoposide or teniposide.
- 7. A pharmaceutical composition according to claim 1, wherein said composition is useful in the treatment of neoplastic diseases.
- 8. A pharmaceutical composition according to claim 1, wherein the 20(S)-camptothecin is 9-amino-20(S)-camptothecin.

10.

- 9. A pharmaceutical composition according to claim 1, wherein the 20(S)-camptothecin is 9-nitro-20(S)-camptothecin.
- delivering to a patient in need of treatment a therapeutically effective amount of 20(S)-camptothecin, an analog of 20(S)camptothecin, or a derivative of 20(S)-camptothecin in combination with an effective amount of at least one of an alkylating agent,

epidophyllotoxin, antimetabolite, antibiotic or vinca alkaloid.

A method for treating disease comprising:

- 11. A method according to claim 10, wherein the alkylating agent is cyclophosphamide, ifosfamide, melphalan, hexamethylmelamine, thiotepa or dacarbazine.
- 12. A method according to claim 10, wherein the antimetabolite is 5-fluorouracil, cytarabine or a folic acid analog.
- 13. A method according to claim 10, wherein the antibiotic is daunorubicine, doxorubicin, bleomycin or mitomycin.
- 14. A method according to claim 10, wherein the vinca alkaloid is vinblastine, vincristine, or their synthetic analogues.
- 15. A method according to claim 10, wherein the epidophyllotoxin is etoposide or teniposide.
- 16. A method according to claim 10, wherein said composition is useful in the treatment of neoplastic diseases.

5

- 17. A method according to claim 10. wherein the 20(S)-camptothecin is 9-amino-20(S)-camptothecin.
- 18. A method according to claim 10, wherein the 20(S)-camptothecin is 9-nitro-20(S)-camptothecin.